

## Clopidogrel (Plavix®) Criteria for Use in Veteran Patients

VHA Pharmacy Benefits Management Strategic Healthcare Group and the Medical Advisory Panel

*The following recommendations are based on current medical evidence. The content of the document is dynamic and will be revised as new clinical data becomes available. The purpose of this document is to assist practitioners in clinical decision-making, to standardize and improve the quality of patient care, and to promote cost-effective drug prescribing. The clinician, however, must make the ultimate judgment regarding the propriety of any course of treatment in light of individual patient situations.*

**The following document briefly outlines recommendations for clopidogrel for various indications and clinical situations, as outlined below. These are not meant to be all inclusive. For other clinical situations and for those with insufficient evidence to determine risk and benefit, consultation with local experts is advised.**

### **Recommendations for use of clopidogrel post percutaneous coronary intervention (PCI)/stent<sup>1,5, 22-33</sup>**

- Patients who receive coronary stents will require dual anti-platelet therapy with aspirin and clopidogrel in order to maintain patency of the artery.
- Clopidogrel should be initiated at the time of the procedure, if not before and ideally continued for 12 months. American Heart Association (AHA) guidelines indicate clopidogrel duration should be guided by stent type [bare metal 1 month, sirolimus (Cypher(r), Cordis) 3 months and paclitaxel (Taxus(r), Boston Scientific) 6 months]. (Level A)
- There is recent evidence that drug eluting stents (DES) may confer a rare but increased risk of late stent thrombosis in certain patient populations (including left main DES, DES of bifurcations, overlapping DES, prior history of late stent thrombosis, and DES of bypass graft), suggesting a longer duration of dual therapy may be needed. The exact duration of clopidogrel therapy in these situations has not been conclusively established, however when making this determination providers must weigh the benefit of prolonged or indefinite clopidogrel therapy against the risk for bleeding on a patient by patient basis. (Level C)<sup>22-32</sup>
- Data emerging from pooled meta-analyses and registries suggest the need for uninterrupted dual antiplatelet therapy (aspirin 325 mg, clopidogrel 75 mg daily) throughout the post-stenting treatment period. Any elective procedures which would require stopping or interrupting this therapy (dental work, colonoscopy, etc.) should be delayed until the minimum treatment duration based on stent type is completed.<sup>33</sup>
- The dose of aspirin following a PCI/stent is 325mg daily. This higher dose of aspirin should be given for one month following bare metal stents, 3 months following sirolimus (Cypher(r), Cordis) stenting, and 6 months post paclitaxel (Taxus(r), Boston Scientific) stents, after which aspirin should be continued indefinitely at a dose of 75 to 162 mg daily.
- Patients undergoing brachytherapy should receive clopidogrel 75 mg and aspirin 325 mg daily for an indefinite period unless there is a bleeding risk. (Level C)

### **Recommendations for use of clopidogrel in NSTEMI/Unstable angina acute coronary syndromes<sup>1,3,5,6,9,11,13</sup>**

- In patients with acute coronary syndrome and /or unstable angina in whom no revascularization procedure is planned, clopidogrel should be added to aspirin as soon as possible for at least 1 month (Level A) and up to 12 months (Level B).
- Clopidogrel will be discontinued after 12 months unless compelling clinical reasons exist to maintain therapy for a longer period (Refer to the section on PCI/Stent for specific guidance on drug eluting stents {DES}).

### **Recommendations for use of clopidogrel in STEMI acute coronary syndrome<sup>18-20</sup>**

- In patients with STEMI < 75 years of age receiving fibrinolytics clopidogrel should be administered as a 300mg loading dose followed by 75mg once daily until hospital discharge (up to 8 days) or longer if undergoing angiography/coronary intervention as described below in recommendations post PCI/Stent. (Level A)
- In patients of any age with STEMI regardless of whether fibrinolytics are utilized clopidogrel should be administered at a dose of 75mg once daily until hospital discharge or up to 4 weeks or longer if undergoing angiography/coronary intervention as described below in recommendations post PCI/Stent. (Level A)
- In patients with STEMI who do not undergo PCI consideration can be given to continuing clopidogrel 75mg once daily for up to 1-year based on extrapolation from trials in NSTEMI/Unstable angina. (Level B)

**Recommendations for use of clopidogrel in stable coronary artery disease (CAD)<sup>17</sup>**

- There is insufficient evidence to recommend initiation in those patients with stable CAD and who do not meet criteria in this document.
- It is recommended that clopidogrel not be added to aspirin therapy for stable/asymptomatic atherosclerosis and/or for patients with multiple cardiovascular risks unless there are other criteria that have been met. (Level A)

**Recommendations for use of clopidogrel in aspirin adverse events<sup>16</sup>**

- Clopidogrel should be used in patients who are aspirin allergic, i.e.; anaphylaxis, aspirin induced bronchospasm. (Level A)
- In patients with a history of gastrointestinal complications from aspirin (i.e.; bleeding, stomach upset), adding a proton pump inhibitor to aspirin therapy is preferred over switching to clopidogrel. (Level B)

**Recommendations for use of clopidogrel for Coronary Artery Bypass Grafting (CABG)<sup>1,14</sup>**

- In patients receiving clopidogrel for ACS, discontinuation is suggested 3-5 days prior to elective CABG. (Level B) and restarted as soon as patient's clinical condition tolerates for up to 12 months.
- For those patients undergoing saphenous vein bypass graft and allergic to aspirin, clopidogrel, 300mg, as a loading dose 6 hours after the operation followed by 75mg daily may be initiated. (Level B).
- Clopidogrel may be continued for 1-3 months post CABG. (Level C)

**Recommendations for the use of clopidogrel in Peripheral Vascular Disease (PVD)<sup>2,4,12</sup>**

- Clopidogrel is not recommended for PVD except in cases of aspirin allergy. (Level B)

**Recommendations for the use of clopidogrel in recurrent cerebral ischemic events<sup>7,10,21</sup>**

- Patients with recurrence of cerebral ischemic events while on therapy with aspirin should be changed to extended release dipyridamole/aspirin. (Level B)
- Clopidogrel is an alternative for those patients who have had recurrent cerebrovascular events, who have a documented aspirin allergy, as mentioned above or are intolerant of extended release dipyridamole (recurrent headache). (Level B)
- The combination of aspirin and clopidogrel is not advised for secondary stroke prophylaxis due to increased risk of adverse events demonstrated in the MATCH trial. (Level B)

**Recommendations for the use of clopidogrel in noncardiac stenting<sup>15</sup>**

- Patients who undergo carotid artery stenting may be initiated on clopidogrel 75 mg/day and continued for 4-6 weeks post stent. (Level B)
- Patients who undergo intracranial stents may require longer durations of therapy and can be continued up to 1 year. (Level B)
- Patients who undergo renal artery stenting may be initiated on clopidogrel 75 mg daily for up to 12 months post intervention. (Level I)
- Patients who undergo other peripheral stents (inguinal, popliteal, etc) may be continued on clopidogrel 75 mg daily for 30 days post intervention. (Level C)

**Recommendation Grading**

<b>A</b>	A strong recommendation that the intervention is always indicated and acceptable
<b>B</b>	A recommendation that the intervention may be useful/effective
<b>C</b>	A recommendation that the intervention may be considered
<b>D</b>	A Recommend that a procedure may be considered not useful / effective, or may be harmful.
<b>I</b>	Insufficient evidence to recommend for or against – the clinician will use their clinical judgment

**References**

1. Smith SC, Feldman TE, Hirshfeld JW, et al. ACC/AHA/SCAI 2005 Guideline Update for Percutaneous Coronary Intervention-Summary Article. Accessed January 17, 2006 [http://www.acc.org/clinical/guidelines/percutaneous/update/index\\_rev.pdf](http://www.acc.org/clinical/guidelines/percutaneous/update/index_rev.pdf)
2. Hirsch AT, Haskal ZJ, Hertzner NR, et al. ACC/AHA Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric and

- abdominal aorta). Accessed January 17, 2006  
<http://www.acc.org/clinical/guidelines/pad/index.pdf>
3. Yusuf S, Mehta SR, Zhao F, et al. Early and late effects of clopidogrel in patients with acute coronary syndrome. *Circulation*. 2003;107:966-972.
  4. Clagett GP, Sobel M, Jackson MR, et al. Antithrombotic therapy in peripheral arterial occlusive disease: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*. 2004 Sep;126(3 Suppl):609S-626S.
  5. Popma JJ, Berger P, Ohman EM, et al. Antithrombotic therapy during percutaneous coronary intervention: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*. 2004 Sep;126(3 Suppl):576S-599S.
  6. Harrington RA, Becker RC, Ezekowitz M, et al. Antithrombotic therapy for coronary artery disease: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*. 2004 Sep;126(3 Suppl):513S-548S.
  7. Albers GW, Amarencu P, Easton JD, Sacco RL, Teal P. Antithrombotic and thrombolytic therapy for ischemic stroke: the Seventh ACCP Conference on Antithrombotic and Thrombolytic Therapy. *Chest*. 2004 Sep;126(3 Suppl):483S-512S.
  8. Weintraub WS, Mahoney EM, Lamy A, et al; CURE Study Investigators. Long-term cost-effectiveness of clopidogrel given for up to one year in patients with acute coronary syndromes without ST-segment elevation. *J Am Coll Cardiol*. 2005 Mar 15;45(6):838-45.
  9. Sabatine MS, Cannon CP, Gibson CM, et al; CLARITY-TIMI 28 Investigators. Addition of clopidogrel to aspirin and fibrinolytic therapy for myocardial infarction with ST-segment elevation. *N Engl J Med*. 2005 Mar 24;352(12):1179-89.
  10. Diener HC, Bogousslavsky J, Brass LM, et al; MATCH investigators. Aspirin and clopidogrel compared with clopidogrel alone after recent ischemic stroke or transient ischemic attack in high-risk patients (MATCH): randomized, double-blind, placebo-controlled trial. *Lancet*. 2004 Jul 24-30;364(9431):331-7.
  11. Kastrati A, von Beckerath N, Joost A, et al. Loading with 600 mg clopidogrel in patients with coronary artery disease with and without chronic clopidogrel therapy. *Circulation*. 2004 Oct 5;110(14):1916-9.
  12. Jagroop IA, Matsagas MI, Geroulakos G, Mikhailidis DP. The effect of clopidogrel, aspirin and both antiplatelet drugs on platelet function in patients with peripheral arterial disease. *Platelets*. 2004 Mar;15(2):117-25.
  13. Patti G, Giuseppe C, Pasceri V et al. Randomized Trial of High Loading Dos of Clopidogrel for Reduction of Periprocedural Myocardial Infarction in Patients Undergoing Coronary Intervention. Results from the ARMYDA-2 (Antiplatelet therapy for Reduction of Myocardial Damage during Angioplasty) Study. *Circulation* 2005;111:2099-2106.
  14. Cannon CP, Mehta SR, Aranki SF. Balancing the Benefit and Risk of Oral Antiplatelet Agents in Coronary Artery Bypass Surgery. *Ann Thorac Surg* 2005;80:768-79.
  15. Payne DA, Hayes CI, Hayes PD et al. Beneficial Effects of Clopidogrel Combined with Aspirin in Reducing Cerebral Emboli in Patients Undergoing Carotid Endarterectomy. *Circulation* 2004; 109(12): 1476-1481.
  16. Chan FKL, Ching JYL, Hung LCT, et al. Clopidogrel versus aspirin and esomeprazole to prevent recurrent ulcer bleeding. *N Engl J Med* 2005;352:238-44.
  17. Bhatt DL, Fox KA, Hacke W, et al. Clopidogrel and Aspirin versus Aspirin Alone for the Prevention of Atherothrombotic Events. *N Engl J Med*. 2006 Mar 12; [Epub ahead of print]
  18. Hahn SA, Chandler C. Diagnosis and management of ST elevation myocardial infarction: a review of the recent literature and practice guidelines. *Mt Sinai J Med*. 2006 Jan;73(1):469-81
  19. Sabatine MS, Morrow DA, Montalescot G, et al; Clopidogrel as Adjunctive Reperfusion Therapy (CLARITY)-Thrombolysis in Myocardial Infarction (TIMI) 28 Investigators. Angiographic and clinical outcomes in patients receiving low-molecular-weight heparin versus unfractionated heparin in ST-elevation myocardial infarction treated with fibrinolytics in the CLARITY-TIMI 28 Trial. *Circulation*. 2005 Dec 20;112(25):3846-54. Epub 2005 Nov 15.
  20. Sabatine MS, Cannon CP, Gibson CM, et al; Clopidogrel as Adjunctive Reperfusion Therapy (CLARITY)-Thrombolysis in Myocardial Infarction (TIMI) 28 Investigators. Effect of clopidogrel pretreatment before percutaneous coronary intervention in patients with ST-elevation myocardial infarction treated with fibrinolytics: the PCI-CLARITY study. *AMA*. 2005 Sep 14;294(10):1224-32.

21. Sacco RL, Adams R, Albers G, et al; American Heart Association/American Stroke Association Council on Stroke; Council on Cardiovascular Radiology and Intervention; American Academy of Neurology. Guidelines for prevention of stroke in patients with ischemic stroke or transient ischemic attack: a statement for healthcare professionals from the American Heart Association/American Stroke Association Council on Stroke:co-sponsored by the Council on Cardiovascular Radiology and Intervention: the American Academy of Neurology affirms the value of this guideline. *Circulation*. 2006 Mar 14;113(10):e409-49.
22. Park DW, Park SW, Park KH, et al. Frequency of and risk factors for stent thrombosis after drug-eluting stent implantation during long-term follow-up. *Am J Cardiol*. 2006 Aug 1;98(3):352-6.
23. Buhk JH, Wellmer A, Knauth M. Late in-stent thrombosis following carotid angioplasty and stenting. *Neurology*. 2006 May 23;66(10):1594-6.
24. Stone GW, Aronow HD. Long-term care after percutaneous coronary intervention: focus on the role of antiplatelet therapy. *Mayo Clin Proc*. 2006 May;81(5):641-52.
25. Urban P, Gershlick AH, Guagliumi G, e-Cypher Investigators. Safety of coronary sirolimus-eluting stents in daily clinical practice: one-year follow-up of the e-Cypher registry. *Circulation*. 2006 Mar 21;113(11):1434-41.
26. Ong AT, McFadden EP, Regar E, et al. Late angiographic stent thrombosis (LAST) events with drug-eluting stents. *J Am Coll Cardiol*. 2005 Jun 21;45(12):2088-92.
27. Iakovou I, Schmidt T, Bonizzoni E, et al. Incidence, predictors, and outcome of thrombosis after successful implantation of drug-eluting stents. *JAMA*. 2005 May 4;293(17):2126-30.
28. Zimarino M, Renda G, De Caterina R. Optimal duration of antiplatelet therapy in recipients of coronary drug-eluting stents. *Drugs*. 2005;65(6):725-32.
29. Spertus JA, Kettelkamp R, Vance C, et al. Prevalence, predictors, and outcomes of premature discontinuation of thienopyridine therapy after drug-eluting stent placement: results from the PREMIER registry. *Circulation*. 2006 Jun 20;113(24):2803-9.
30. Pfisterer ME. Basel Stent Cost-effectiveness Trial-Late Thrombotic Events (BASKET LATE) Trial. Presented at 55th Annual Scientific Session The American College of Cardiology Scientific Session. Atlanta, GA. 2006
31. Bavry AA, Kumbhani DJ, Helton TJ, Borek PP, Mood GR, Bhatt DL. Late thrombosis of drug-eluting stents: a meta-analysis of randomized clinical trials. *Am J Med*. 2006 Dec;119(12):1056-61.
32. Eisenstein EL, Anstrom KJ, Kong DF, et al. Clopidogrel Use and Long-term Clinical Outcomes After Drug-Eluting Stent Implantation. *JAMA*. 2007;297(2):159-68.
33. Grines CL, Bonow RO, Casey DE Jr; et al. Prevention of Premature Discontinuation of Dual Antiplatelet Therapy in Patients With Coronary Artery Stents. A Science Advisory From the American Heart Association, American College of Cardiology, Society for Cardiovascular Angiography and Interventions, American College of Surgeons, and American Dental Association, With Representation From the American College of Physicians. *Circulation*. 2007 Jan 15; [Epub ahead of print]

### Criteria Checklist for Clopidogrel

Pharmacy Benefits Management Strategic Healthcare Group and the Medical Advisory Panel

<b>Indication for therapy</b>	<b>Response</b>
Patient with one of the following conditions: <input type="checkbox"/> unstable angina <input type="checkbox"/> Acute coronary syndrome( NSTEMI or STEMI) <input type="checkbox"/> Post PCI/stent placement <input type="checkbox"/> Documented aspirin allergy requiring antiplatelet therapy <input type="checkbox"/> Intracranial/carotid stent <input type="checkbox"/> Brachytherapy <input type="checkbox"/> Peripheral stent (renal, inguinal, popliteal) <input type="checkbox"/> Cerebrovascular disease with recurrent ischemia	<input type="checkbox"/> yes <input type="checkbox"/> no <b><i>If no, patient is ineligible to receive clopidogrel</i></b>
<b>Duration of therapy</b> <input type="checkbox"/> 1 month-post stent placement <input type="checkbox"/> 3 months- post ACS <input type="checkbox"/> 9 months- high risk post ACS <input type="checkbox"/> 12 month- post stent placement <input type="checkbox"/> indefinite- aspirin allergic patient <input type="checkbox"/> peripheral stent duration_____ months <input type="checkbox"/> other _____	
<b>Dosing</b> <input type="checkbox"/> ACS- 300 mg load orally then daily dose of 75 mg <input type="checkbox"/> ACS-600 mg load orally than daily dose of 75 mg <input type="checkbox"/> Post stent placement-300 mg load orally then daily dose of 75 mg <input type="checkbox"/> Non acute conditions- 75 mg daily	
<b>Monitoring</b> <input type="checkbox"/> Out-patients should have a cardiology referral for indefinite ACS use or longer than 3 month occurrence of ACS <input type="checkbox"/> Patients with cerebrovascular disease should have a neurology referral if clopidogrel and aspirin are used together <input type="checkbox"/> Patients should be followed for development of neutropenia or thrombotic thrombocytopenic purpura	
<b>Contraindications</b> Any of the following: <input type="checkbox"/> Active bleeding <input type="checkbox"/> History of bleeding diathesis <input type="checkbox"/> Known hypersensitivity to clopidogrel or any component of the product	<input type="checkbox"/> yes <input type="checkbox"/> no <b><i>If yes, patient is ineligible to receive clopidogrel</i></b>